

Click here for original study (Below)

## **High-Intensity Exercise on the ROM Training Device Improves VO2 Max and Endurance Capacity in Untrained Adults**

Eline M. van Es, Hanno van der Loo. TNO Defence, Security and Safety, Soesterberg, The Netherlands

Email: [Eline.vanes@tno.nl](mailto:Eline.vanes@tno.nl)

*Official Journal of the American College of Sports Medicine Vol. 39 No.5 Supplement S349 – 2005 (May 2007)*

The importance of sports and physical activity for the maintenance and improvement of health and physical fitness is increasingly recognized. However, most people do not allow themselves the time to engage in a conventional, high volume – moderate intensity training program. A less time consuming, yet equally effective program could solve this problem.

**Purpose:** To investigate whether the VO2 Max and endurance capacity of healthy, untrained adults can be improved by a high-intensity – low-volume exercise program using a whole body training device, the so-called ROM machine.

**Methods:** 16 healthy, untrained subjects (12 men, 4 women, age  $36 \pm 10$  years) exercised for 8 weeks, 3 times a week. Net exercise time per session was 8 minutes. In every session, first the upper body was exercised for 4 minutes and then, after a short break, the lower body was exercised for the same length of time. The upper body exercise is more or less similar to rowing, with resistance in both movement directions. The lower body exercise can be described as deep stepping. Subjects were encouraged to train at their maximal intensity. Total net exercise time over 8 weeks was 3 hours and 12 minutes. Before and after the training period, two tests on a cycle ergometer were conducted: one progressive test to determine VO2 max, maximal power output (Wmax) and maximal heart rate (HRmax) and one constant power test until exhaustion at 80% VO2 max to determine endurance capacity.


**Results:** Both VO2max ( $39.6 \pm 4.7$  vs.  $41.0 \pm 4.8$  ml/kg\*min,  $p=0.01$ ) and Wmax ( $248 \pm 64$  vs.  $272 \pm 66$ W,  $p<0.01$ ) increased significantly over the training period, while HRmax ( $182 \pm 8$  vs.  $181 \pm 8$  bpm) did not change. There was a significant increase of 72% in endurance time at 80% VO2max ( $14:51 \pm 6:31$  vs.  $25:31 \pm 11.57$  min,  $p<0.01$ ).

**Conclusion:** Whole body, high-intensity training on the ROM training device is a time efficient way to improve VO2max and especially endurance capacity in healthy, untrained adults.



Eline M. van Es  
 Hanno van der Loo  
 TNO Defence, Security and Safety, Soesterberg, the Netherlands.

Email: Eline.vanes@tno.nl



# High-intensity exercise on the ROM training device improves VO<sub>2</sub>max and endurance capacity in untrained adults

## INTRODUCTION

The importance of sports and physical activity for the maintenance and improvement of health and physical fitness is increasingly recognized. However, most people do not allow themselves the time to engage in a conventional, high volume - moderate intensity training program. A less time consuming, yet equally effective program could solve this problem. This study investigated whether the physical fitness of healthy, untrained volunteers in the age group 20-55 years can be improved by a specific kind of high intensity training: the so called Fast Fit concept. The training program consisted of three eight-minute exercise sessions per week on the so called ROM training device.




## METHODS

Twelve men and four women (age 36 ± 10 years) trained for eight weeks, three times a week. In every session, first the upper body was exercised for four minutes and then, after a short break, the lower body was exercised for the same length of time. The upper body exercise is more or less similar to rowing, with resistance in both movement directions. The lower body exercise can be described as deep stepping. Subjects were encouraged to train at their maximal intensity. Total net exercise time over eight weeks was 3 hours and 12 minutes. Before and after the training period, two tests on a cycle ergometer were conducted; one progressive test to determine maximal oxygen uptake (VO<sub>2</sub>max), maximal power output (Wmax) and maximal heart rate (HRmax) and one constant power test until exhaustion at 80% VO<sub>2</sub>max to determine endurance capacity.

## RESULTS

The training induced (see table):  
 • a significant increase of VO<sub>2</sub>max;  
 • a significant increase of the endurance time in cycling at 80% VO<sub>2</sub>max (see fig);  
 • a significant increase of maximal external power during a progressive cycling test.  
 No injuries or other adverse effects were recorded.

	Pre-test	Post-test	Relative change	p-value
VO <sub>2</sub> max (ml/kg/min)	38.6 ± 6.7	49.9 ± 8.8	+29.5%	<0.001*
Endurance time 80% VO <sub>2</sub> max (min)	16:00 ± 0:15	18:52 ± 0:18	+17.5%	<0.01*
Maximal power output (W)	308 ± 64	372 ± 85	+20.7%	<0.05*
HRmax (b/min)	173 ± 8.36	173 ± 8.07	0%	>0.05
Maximal heart rate (b/min)	182 ± 9	181 ± 8	-0.5%	>0.05
Endurance time 80% VO <sub>2</sub> max (min)	14:51 ± 0:31	20:21 ± 1:07	+35.5%	<0.001*
Mean VO <sub>2</sub> 80% VO <sub>2</sub> max test (ml/kg/min)	33.2 ± 4.7	33.4 ± 4.1	+0.3%	>0.05
Mean HR 80% VO <sub>2</sub> max test (b/min)	170 ± 8.06	176 ± 8.04	+3.2%	<0.05*



## DISCUSSION

The effect of training depends on the intensity, duration and frequency of training and the length of the training period. In the table below, the training effects in this study are compared to some conventional training schedules. Eight weeks of training on the ROM training device is very time efficient in improving VO<sub>2</sub>max. However, the improvement per week of training is more pronounced in training programs with more training time.


Study	Training weeks	Training time (min)	VO <sub>2</sub> max increase (%)	Increase per week (%)	Minutes training per 1% increase
Wagner et al (1990)	12	1425	4.3	0.36	400
Wolfe et al (1990)	7	1425	14.3	2.05	210
Tabata et al (1996)	6	1800	9.4	1.57	150
Wagner et al (1990)	8	720	7.4	0.93	97
E.M. van Es et al (2006)	8	180	3.3	0.41	35

Both the increase in VO<sub>2</sub>max and the increase in endurance time at 80% VO<sub>2</sub>max are comparable with the effects found in other studies on high intensity training (Gorostiaga et al, 1991; Tabata et al, 1996; Burgomaster et al, 2005; Franch et al, 1998).

## CONCLUSION

Whole body, high-intensity training on the ROM training device is a time efficient way to improve VO<sub>2</sub>max and especially endurance capacity in healthy, untrained adults.

*Acknowledgements:*  
 This work was funded by Fast Fit BV, Zutphen, The Netherlands. We are indebted to our colleague Ineke Klöpping, Maikel Rondon (Fast Fit), Boris Sala (Exercise Worx) and to our subjects.



References:  
 Burgomaster KA, Hughes SC, Heigenhauser GJ, Bradwell SN & Gibala MJ (2005). Six sessions of sprint interval training increases muscle oxidative potential and cycle endurance capacity in humans. *Journal of Applied Physiology*, 98: 1985-1990.  
 E.M. van Es et al (2006). Efficiency of physical fitness training according to the Fast Fit concept. TNO Defence, Security and Safety report DV 2006 C299.  
 Franch J, Madson R, Djurhuus MS, Pedersen PK (1998). Improved running economy following intensified training correlates with reduced ventilatory demands. *Medicine and Science in Sports and Exercise*, 30: 1250-1256.  
 Gorostiaga EM, Whaley CB, Foster C & Hickson BC (1991). Uniqueness of interval and continuous training at the same maintained exercise intensity. *European Journal of Applied Physiology and Occupational Physiology*, 63: 101-107.  
 McKenzie S, Phillips SM, Carter SL, Lowther L, Gibala MJ & Tarnopolsky MA (2000). Endurance exercise training attenuates leucine oxidation and BCOAD activation during exercise in humans. *American Journal of Physiology, Endocrinology and Metabolism*, 278: S80-S87.  
 Suzuki S, Umeta G, Ishida Y, Kawahira H & Yamamura M (1998). Influences of low intensity exercise on body composition, food intake and aerobic power of sedentary young females. *Applied Human Science*, 17: 259-266.  
 Tabata I, Nishimura K, Knuzaki M, Hirai Y, Ogita R, Miyachi M & Yamamoto K (1996). Effects of moderate-intensity endurance and high-intensity intermittent training on anaerobic capacity and VO<sub>2</sub>max. *Medicine and Science in Sports and Exercise*, 28: 1327-1330.